



AC JOINT INJURIES

EXPERT CONSULTANT | Edward McFarland, MD

What is the AC joint?

The AC (acromioclavicular) joint is a joint in the **shoulder** where the collarbone (clavicle) meets the shoulder blade (scapula). The specific part of the scapula adjacent to the clavicle is called the acromion, hence the name AC joint.

What kinds of problems occur at the AC joint?

The most common problems that occur at the AC joint are arthritis, fractures and “separations.” Arthritis is a condition characterized by loss of cartilage in the joint. AC joint arthritis is common in weight lifters, especially with the bench press, and (to a lesser

extent) military press. AC joint arthritis may also be present when there are rotator cuff problems.

What is an AC separation?

When the AC joint is “separated,” it means that the ligaments connecting the acromion and clavicle have been damaged, and the two structures no longer line up correctly. AC separations can be anywhere from mild to severe, and AC separations are “graded” depending upon which ligaments are torn and how badly they are torn.

Grade I Injury — the least damage is done, and the AC joint still lines up.

Grade II Injury — damage to the ligaments which reinforce the AC joint. In a grade II injury these ligaments are only stretched but not entirely torn. When stressed, the AC joint becomes painful and unstable.

Grade III Injury — AC and secondary ligaments are completely torn and the collarbone is no longer tethered to the shoulder blade, resulting in a visible deformity.

How is AC joint arthritis treated?


If rest, ice, medication, and modifying activity does not work, then the next step is a cortisone shot. One shot into the joint sometimes takes care of the pain and swelling permanently, although the

Statistics

- ▶ In 2008, nearly 95,000 people visited a physician with an AC joint related injury.
- ▶ More males than females suffer AC joint injuries.

Athletes who have had AC joint injuries

- ▶ Sam Bradford, St. Louis Rams Quarterback
- ▶ Matthew Stafford, Detroit Lions



effect is unpredictable and may not be permanent. **Surgery** may be indicated if nonsurgical measures fail. Since the pain is due to the ends of the bones making contact with each other, the treatment is removal of a portion of the end of the clavicle. This outpatient surgery can be performed through a small incision about one inch long or arthroscopically using several small incisions. Regardless of the technique utilized, the recovery and results are about the same. Most patients have full motion by six weeks and return to sports by 12 weeks.

What is the treatment for AC separation?

These can be very painful injuries, so the initial treatment is to decrease pain. This is best accomplished by immobilizing the arm in a sling, and placing an ice pack to the shoulder for 20–30 minutes every two hours as needed.



AC JOINT INJURIES

Acetaminophen and non-steroidal anti-inflammatory drugs can also help the pain. As the pain starts to subside, it is important to begin moving the fingers, wrist, and elbow, and eventually the shoulder in order to prevent a stiff or “frozen” shoulder. The length of time needed to regain full motion and function depends on the severity or grade of the injury. Recovery from a Grade I AC separation usually takes 10 to 14 days, whereas a Grade III may take six to eight weeks.

When is surgery indicated?

Grade I and II separations very rarely require surgery. Even Grade III injuries usually allow return to full activity with few restrictions. In some cases a painful lump may persist, necessitating partial clavicle excision in selected individuals such as high caliber throwing athletes. Surgery can be very successful in these cases, but as always, the benefits must be weighed against the potential risks.

Additional Information

www.sportsmed.org

www.orthoinfo.org

www.STOPSportsInjuries.org